

Seed dispersal of *Attalea phalerata* (Palmae) by Crested caracaras (*Caracara plancus*) in the Pantanal and a review of frugivory by raptors

Mauro Galetti¹ and Paulo R. Guimarães Jr.^{1,2}

¹ Plant Phenology and Seed Dispersal Research Group, Departamento de Ecologia, Universidade Estadual Paulista (UNESP), Caixa Postal 199, 13506-900 Rio Claro, São Paulo, Brazil, mgaletti@rc.unesp.br

² Programa de Pós-Graduação em Ecologia, Instituto de Biologia, Universidade Estadual de Campinas, Caixa Postal 6109, 13083-970, Campinas, SP, Brazil. paulomiudo@uol.com.br

Recebido em 26 de agosto de 2004; aceito em 20 de outubro de 2004

ABSTRACT: We observed Crested caracaras (*Caracara plancus*) consuming and dispersing fruits of the palm *Attalea phalerata* at Pantanal, Brazil. We reviewed the literature of seed dispersal by raptors and suggest that raptors may affect seed dispersal by three different paths: secondary seed dispersal by preying on frugivorous birds, primary seed dispersal of ornithocoric fruits and primary seed dispersal of large, lipid-rich fruits. The latter path may be an important long-distance seed dispersal mechanism for large seeds.

KEY WORDS: birds of prey, megafauna fruits, *Polyborus*, Pantanal, raptors, secondary seed dispersal, frugivory

Raptors are not the typical image of a frugivore bird and their role in seed dispersal have been neglected in reviews of the importance of birds for plants (Snow 1981, Fleming 1991). However, raptors can be important secondary seed dispersers when they feed on frugivore preys or primary dispersers when they deliberately consume fruits (Engel 2000, Shanahan *et al.* 2001, Nogales *et al.* 2002). Available information on raptor seed dispersal is scanty and based mainly on anecdotal records. Here we report the seed dispersal of the “bacuri” palm *Attalea phalerata* (Palmae) by Crested caracaras *Caracara plancus* (Falconidae) in the Pantanal of Brazil and review the evidences of seed dispersal by raptors around the world.

Crested caracaras are generalist falconids that are known to feeding on a variety of food items, including carrion, small vertebrates such as lizards, frogs and rodents, insects, eggs and fruits (Sick 1997). Although widespread in Pantanal and other natural (or human-generated) open fields, few published studies focus on the ecology of this species. On July and November 2003, we regularly observed interactions of Crested caracaras and *A. phalerata* fruits. *Attalea phalerata* produces large-sized fruits, rich in lipids (da Silva *et al.* 1994), that are dispersed by a small coterie of animals, including two introduced species (cattle, and feral pigs), and rheas, tapirs, and rodents (agoutis and *Clyomys* spp.) (M. Galetti unpublished data). Rodents are mainly seed predators but also disperse *Attalea* seeds at short distance, while tapirs, cattle and feral pigs can be long distance seed dispersers (Fragoso *et al.* 2003). Hyacinth macaws are the main seed predator and also carry *Attalea* nuts and may occasionally disperse its seeds (Guedes and Harper 1995). Of the recorded *Caracara-Attalea* interactions, in two occasions we observed the cacaracas taking off with one *Attalea* fruit in its beak (Figure 1) and perching in trees more than 200 m away. Caracaras eat only the pulp of the fruit, leaving the nut intact.

We also compiled the evidence of frugivory and seed dispersal by four families of birds of prey (Accipitridae,



Figure 1. *Caracara plancus* dispersing seeds of *Attalea phalerata* (Palmae) at Fazenda Rio Negro, Pantanal, Brazil.

Cathartidae, Falconidae, and Pandionidae) using the ISI-Web of Science (isi3.isiknowledge.com), the Raptor Information System (ris.wr.usgs.gov), and different books about natural history of birds (see Table 1). We recorded evidence of frugivory in 13 taxa of birds of prey (three families, Table 1). The records on frugivory by raptors are scarce, but sampled data indicate three paths in which birds of prey may disperse seeds.

First, raptors may act as secondary seed dispersers by consuming primary seed dispersers and seed predators and discarding intact seeds at the perch or regurgitating pellets with seeds (Hall 1987, Dean and Milton 1988, Engel 2000, Pearson and Ortega 2001, Shanahan *et al.* 2001). The strongest evidence of the ecological importance of this kind of dispersal is restricted to a single kite species (*Falco tinnunculus*) and non-falconiform predatory birds in ocean islands (Nogales *et al.* 1998, 2002), but there is evidence of its occurrence for introduced predatory mammals, such as

Table 1. Evidence of frugivory by birds of prey (Cathartidae, Accipitridae, Falconidae).

Order	Family	Species	Fruits recorded ¹	Distribution ²	Reference	
Ciconiformes	Cathartidae	<i>Cathartes aura</i>	<i>Acrocomia</i> (P)*, <i>Elaeis</i> (P)*, <i>Juniperus</i> (C), grapes	NA	Hiraldo et al. 1991, Sick 1997	
		<i>Coragyps atratus</i>	“fruit”	NA	Buskirk & Lechner 1978	
Falconiformes	Accipitridae	<i>Buteogallus urubitinga</i>	<i>Spondias</i> (A)*	NA	Del Hoyo et al. 1995, Sick 1997	
		<i>Elanoides forficatus</i>	<i>Byrsonima</i> (Ma), <i>Castilla</i> (Mo), <i>Cupania</i> (S), <i>Matayba</i> (S), <i>Sapium</i> (E), “fruits” (S and Mo)	NA	Buskirk & Lechner 1978, Lemke 1979, Del Hoyo et al. 1995, Sick 1997, Chaves 2000, Gerhardt et al. 2004, Meyer et al. 2004,	
		<i>Gypohierax angolensis</i>	<i>Elaeis</i> (P)*, <i>Raphia</i> (P)*, <i>Phoenix</i> (P)*	AFR	Brown & Amadon 1968, Brown et al. 1982, Del Hoyo et al. 1995	
		<i>Milvus migrans</i>	<i>Elaeis</i> (P)*	Palaearctic	Brown et al. 1982, Del Hoyo et al. 1995	
		<i>Polyboroides typus</i>	<i>Elaeis</i> (P)*	AFR	Brown & Amadon 1968, Brown et al. 1982, Del Hoyo et al. 1995	
		Falconidae	<i>Caracara plancus</i>	<i>Attalea</i> (P)*, <i>Elaeis</i> (P)*, <i>Cocos</i> (P)*, “beans” (L), “peanuts” (L),	NA	Del Hoyo et al. 1995, Sick (1997), This work
			<i>Daptrius ater</i>	<i>Elaeis</i> (P)*, <i>Desmoncus</i> (P), <i>Mauritia</i> (P)*, “unidentified seeds”	NA	Schubart et al. 1965, Brown & Amadon 1968, Thiollay 1991, Del Hoyo et al. 1995, Sick 1997
			<i>Daptrius americanus</i>	<i>Poraqueiba</i> (I), “fruits” (Ma and P), “unidentified fruits”	NA	Schubart et al. 1965, Thiollay 1991, Del Hoyo et al. 1995, Sick 1997
			<i>Falco rufigularis</i>	“small green fruit” +	NA	P. Madrigal, pers. comm.
			<i>Milvago chimachima</i>	<i>Elaeis</i> (P)*, <i>Byrsonima</i> (Ma)	NA	Schubart et al. 1965, Del Hoyo et al. 1995, Sick 1997
		<i>Micrastur ruficollis</i>	<i>Blomia</i> (S)	NA	Thorstrom 1996	

¹(A) Anacardiaceae, (C) Cupressaceae, (E) Euphorbiaceae, (I) Icacinaceae, (L) Leguminosae, (Ma) Malpighiaceae, (Mo) Moraceae, (P) Palmae, (S) Sapindaceae, * large fruits (> 3 cm of length), + not confirmed, ²AFR Africa, NA Neotropical America

feral cats (Nogales *et al.* 1996) and for several continental predatory birds (review in Dean & Milton 1988). This path of seed dispersal is more likely to affect seedling recruitment in simple ecological systems, such as islands, in which frugivores consume few plant species and are consumed by few predators in a regular basis (Nogales *et al.* 2002). However, based on the quantity of preys - especially seed-eating rodents- consumed by predatory birds, Dean & Milton (1988) suggest that the amounts of seeds that are secondarily dispersed by predatory birds are not negligible. Differently from other kinds of secondary seed dispersal, such as short distance, scatterhoarding by rodents, secondary dispersal by raptors is likely to enhance considerably seed dispersal distances (Nogales *et al.* 2002). Further work is needed to clarify the role of the secondary dispersal by birds of prey in continental habitats.

Second, raptors may act as primary seed dispersers of ornithocoric fruits. There is evidence of kites that actively

consume small berries and seeds with fleshy arils. The few reports of this kind of seed dispersal by raptors occur in insectivorous species such as *Elanoides forficatus* (by far, the raptor species in which the frugivory was best recorded) and vertebrate predators such as *Micrastur ruficollis* (Table 1). Raptors capture the fruits on trees, as in *E. fortificatus*, or fallen fruits, such as in *M. ruficollis*. Based on the scarcity of natural history data for several forest-dwelling species, it is possible that more species of kites and hawks eventually consume fruits, acting as non-specialized frugivores and, occasionally, dispersing seeds.

Third, raptors may primarily disperse large-seeded fruits. The consumption of large fruits, especially palm fruits, is apparently widespread among the Neotropical Falconidae with a generalist diet that includes carrion, small vertebrates and insects, but also in wasp-specialist species (*Daptrius ater*) (Table 1). Sick (1997) recorded Crested caracaras consuming fruits of *Elaeis oleifera*, but he did not provide any estimative

of the potential role as seed dispersers. Moreover, opportunistic raptors from other families, such as *Cathartes aura* (Cathartidae), and *Gypohierax angolensis* (Accipitridae) also consume large fruits, including palms (Table 1).

All large fruits recorded here (*Acrocomia*, *Attalea*, *Elaeis*, *Phoenix*, *Raphia*, *Mauritia*, and *Spondias*) have pulps rich in lipids (Carciofi 2002, da Silva *et al.* 1994). The exploitation of lipid-rich fruits by carnivorous is a pattern also recorded by carnivorous arthropods, such as Ponerinae ants and harvestmen, in which lipid-rich fruits are used because they are chemically analogous to animal prey (Machado and Pizo 2000, Pizo and Oliveira 2001). We suggest that a similar explanation may be used to fruit consumption by generalist or insect-specialized raptors.

The interactions of large fruits with large, generalist birds of prey possible have consequences for seed dispersal of these plants. Most of large fruits cited in Table 1 fit to "megafauna seed dispersal syndrome", that is, their putative seed dispersers are large mammals (Janzen and Martin 1982, P. R. Guimarães *et al.* unpublished data). All megafauna are now extinct in South America, except for tapirs, and these large-seeded plants are now dispersed by a small coterie of species, mainly by short-distance scatterhoarding rodents such as agoutis and *Clyomys* (M. Galetti and C. Donatti unpublished data). Thus, occasional dispersal by Crested caracaras and other generalist raptors may provide important events of long-distance seed dispersal.

ACKNOWLEDGEMENTS

MG receives support from CNPq (Bolsa de Produtividade), FAPESP and IFS. PRG receive a fellowship from FAPESP. The project at Fazenda Rio Negro was possible due to the support from Earthwatch Institute (and the volunteers), Conservation International and Instituto de Biologia da Conservação. Two anonymous reviewers provided valuable comments on an earlier version of the manuscript. Thanks to L. Chaves, S. Compton, A. Costa, T. Engel, P. Jordano, J. Lisboa, P. Madrigal, S. Milton, M. Shanahan, S. van Noort, and W. R. Silva for help with information and bibliography.

REFERENCES

- Brown, L.H. and D. Amadon. (1968) *Eagles, Hawks & Falcons of the World*. Wellfleet Press, New Jersey.
- Brown, L. H., Urban E. K., and Newman K. (1982) *The Birds of Africa*. Academic Press, London.
- Buskirk, W. H. and Lechner M. (1978) Frugivory by swallow-tailed kites in Costa Rica. *Auk* 95:767-768.
- Carciofi, A. C. (2002) Estudos sobre nutrição de psitacídeos em vida livre: o exemplo da Arara-Azul (*Anodorhynchus hyacinthinus*). *Ecologia e conservação de psitacídeos no Brasil*. M. Galetti and M. A. Pizo. Melopsittacus Publicações Científicas, Belo Horizonte.
- Chaves, L. (2000) Nuevos datos sobre canto, forrejeos y dieta de cuatro especies de aves en Costa Rica. *Zedonia* 4:1.
- da Silva, J.A., da Silva D.B., Junqueira N.T. and Andrade L.R. (1994) *Frutas nativas dos Cerrados*. EMBRAPA, Brasília, DF.
- Dean, W. R. J. and S. J. Milton (1988). Dispersal of seeds by raptors. *African Journal of Ecology* 26: 173-176.
- del Hoyo, J., Elliott A., Sargatal J. and Cabot J. (1992) *Handbook of the birds of the world: New World vultures to guineafowl*, vol. 2. Lynx Edicions, Barcelona, Spain.
- Engel, T. R. (2000) Seed dispersal and forest regeneration in a tropical biocoenosis (Shimba Hills, Kenya. Logos Verlag, Berlin.
- Fleming, T. H. (1991) Fruiting plant - frugivore mutualism: the evolutionary theater and the ecological play. *Plant-animal interactions: evolutionary ecology in tropical and temperate regions*. P. W. Price, T. M. Lewinsohn, G.W. Fernandes and W. W. Benson. John Wiley & Sons: 119-144.
- Fragoso J.M.V., Silvius K.M. and Correa J.A. (2003) Long-distance seed dispersal by tapirs increases seed survival and aggregates tropical trees. *Ecology* 84, 1998-2006
- Gerhardt R. P., Gerhardt D. M., and Vasquez M. A. (2004) Food delivered to nests of Swallow-tailed Kites in Tikal National Park, Guatemala. *Condor* 106:177-181.
- Guedes, N. M. R. and L. H. Harper (1995). Hyacinth macaws in the Pantanal. *The Large Macaws*. J. Abramson, B. L. Speer and J. B. Thomsen. Fort Bragg, Raintree Publications.
- Hall, G. (1987) Seed dispersal by birds of prey. *Zimbabwe Science News* 21:2.
- Hiraldo F., Delibes, M. and Donazar J. A. (1991) Comparison of diets of turkey vultures in 3 regions of northern Mexico. *Journal of Field Ornithology* 319-324.
- Janzen, D. H. and P. S. Martin. (1982) Neotropical Anachonisms: the fruits the gomphotheres ate. *Science* 215:19-27.
- Lemke, T. O. (1979). Fruit-eating behavior of Swallow-tailed kites (*Elanoides forficatus*) in Colombia. *Condor* 81: 207-208.
- Machado, G., and M. A. Pizo. (2000) The use of fruits by the neotropical harvestman *Neosadocus variabilis* (Opiliones, Laniatores, Gonyleptidae). *Journal of Arachnology* 28:357-360.
- Meyer K. D., McGehee S. M., and Collopy M. W. (2004) Food deliveries at Swallow-tailed Kite nests in southern Florida. *Condor* 106:171-176.
- Nogales M., Delgado J.D. and Medina F.M. (1998) Shrikes, lizards and *Lycium intricatum* (Solanaceae) fruits: a case of indirect seed dispersal on an oceanic island (Alegranza, Canary Islands). *Journal of Ecology* 86, 866-871
- Nogales M., Medina F.M. and Valido A. (1996) Indirect seed dispersal by the feral cats *Felis catus* in island ecosystems (Canary Islands). *Ecography* 19, 3-6
- Nogales M., Quilis V., Medina F.M., Mora J.L. and Trigo L.S. (2002) Are predatory birds effective secondary seed dispersers? *Biological Journal of the Linnean Society* 75: 345-352
- Pearson, D. E. and Y. K. Ortega. (2001) Evidence of an indirect dispersal pathway for Spotted Knapweed, *Centaurea maculosa*, seeds, via Deer Mice, *Peromyscus maniculatus*, and Great Horned Owls, *Bubo virginianus*. *Canadian Field Naturalist* 115:354-354.
- Pizo, M. A. and P. S. Oliveira. 2001. Size and lipid content of nonmyrmecochorous diaspores: effects on the interaction with litter-foraging ants in the Atlantic rain forest of Brazil. *Plant Ecology* 157:37-52.
- Shanahan M., So S., Compton S. G., and Corlett R. (2001) Fig-eating by vertebrate frugivores: a global review. *Biological Reviews* 76:529-572
- Schubart, O., Aguirre, A. C., and Sick, H. (1965) Contribuição para o conhecimento da alimentação das aves brasileiras. *Arquivos de Zoologia* 12:95-
- Sick, H. (1997) *Ornitologia Brasileira*. Nova Fronteira, Rio de Janeiro, RJ.
- Snow, D. W. (1981). Tropical frugivorous birds and their food plants: a world survey. *Biotropica* 13: 1-14.
- Thiollay, J. M. (1991) Foraging, home range use and social-behavior of a group-living rain-forest raptor, the red-throated caracara *Daptrius americanus*. *Ibis* 133:382-393.
- Thorstrom, R. (1996) Fruit-eating behavior of a barred forest-falcon. *Journal of Raptor Research* 30:44-44.